



## 7-Substituted Hexahydroquinoline Derivatives and their Calcium Channel Modulator Effects

Miyase G. GÜNDÜZ <sup>1</sup>, Semra ÇELEBİ <sup>2</sup>, Bilgin KAYGISIZ <sup>2</sup>, Rahime ŞİMŞEK <sup>1</sup>\*,  
Kevser EROL <sup>2</sup> & Cihat ŞAFAK <sup>1</sup>

<sup>1</sup> Hacettepe University, Faculty of Pharmacy, Department of Pharmaceutical Chemistry,  
06100, Ankara - Turkey

<sup>2</sup> Osmangazi University, Faculty of Medicine, Department of Pharmacology, Eskisehir-Turkey

---

**SUMMARY.** In this study, calcium antagonistic activity of sixteen hexahydroquinoline derivatives have been elucidated in rat ileum and rat thorasic aorta. In the studies on isolated rat ileum, it has been observed that all compounds showed meaningfull activity. In rat thorasic aorta studies, compound 3a has been found more active than nicardipine.

---

**KEY WORDS:** Calcium antagonistic activity, Hexahydroquinoline, Nicardipine, Synthesis.

\* Author to whom correspondence should be addressed. E-mail: rsimsek@hacettepe.edu.tr

\* Rahime Simsek, PhD